

Human Recombinant CD86 Stable Cell Line

Cat. No.: FCC-137

■ Introduction


Cell Line Name:	CHO-K1/CD86
Gene Synonyms:	B70; B7-2; B7.2; LAB72; CD28LG2
Expressed Gene:	CD86
Host Cell:	CHO-K1, (ATCC® CCL-61™)
Quantity:	Two vials of frozen cells (>1×10 ⁶ cells/vial)
Application:	Binding assay or use as immunogen
Cryopreservation Medium:	70% Complete growth medium, 20% FBS, 10% (V/V) DMSO
Complete Growth Medium:	F12K, 10% FBS
Drug Screening:	12 µg/ml Puromycin
Mycoplasma Contamination:	Not detected
Storage:	Liquid nitrogen immediately upon receipt

■ Background

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.

■ **Representative Data**

Protein Expression Validation



Sample Name	Subset Name	Count
CD86-2.fcs	Lymphocytes	11485
CTRL-CD86.fcs	Lymphocytes	10531

Figure 1: FACS analysis of CD86 expression in CHO-K1/CD86 cells

■ **Cell Culture Procedure**

1. Complete Growth Medium

The base medium for this cell line is ATCC-formulated F-12K Medium, Catalog No. 30-2004. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.

2. Subculturing

- 1) Remove and discard culture medium.
- 2) Briefly rinse the cell layer with 0.25% (w/v) Trypsin- 0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor.
- 3) Add 2.0 mL of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 2 to 5 minutes). **Note:** To avoid clumping do not agitate the cells

by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.

- 4) Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting.
- 5) Add appropriate aliquots of the cell suspension to new culture vessels.
- 6) Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:5 is recommended.

Medium Renewal: Once or twice between subculture.

3. Cryopreservation

Freeze medium: 70% Complete growth medium, 20% FBS, 10% (V/V) DMSO

Storage temperature: liquid nitrogen vapor phase

4. Culture Conditions

Temperature: 37°C,

CO₂ : 5%

■ Limited Use License Agreement

This is a legal agreement between you (Licensee) and Fubio (Suzhou) Biotechnology Co., Ltd governing use of Fubio's stable cell line products and protocols provided to licensee. By purchasing and using the stable cell line, the buyer agrees to comply with the following terms and conditions of this label license and recognizes and agrees to such restrictions:

- 1) The products are not transferable and will be used at the site where they were purchased. Transfer to another site owned by buyer will be permitted only upon written request by buyer followed by subsequent written approval by Fubio.
- 2) The purchaser cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party.
- 3) The products sold by Fubio are for laboratory and animal research purposes only. The products are not to be used on humans, for consumption, or for any unlawful uses.